

Dietary Reference Intakes (DRIs): Tolerable Upper Intake Levels, Vitamins
Food and Nutrition Board, Institute of Medicine, National Academies

Life Stage Group	Vitamin A (µg/d) ^a	Vitamin C (mg/d)	Vitamin D (µg/d)	Vitamin E (mg/d) ^{b,c}	Vitamin K	Thiamin	Riboflavin	Niacin (mg/d) ^c	Vitamin B ₆ (mg/d)	Folate (µg/d) ^c	Vitamin B ₁₂	Pantothenic Acid	Biotin	Choline (g/d)	Carotenoids ^d
Infants															
0 to 6 mo	600	ND ^e	25	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
6 to 12 mo	600	ND	38	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
Children															
1–3 y	600	400	63	200	ND	ND	ND	10	30	300	ND	ND	ND	1.0	ND
4–8 y	900	650	75	300	ND	ND	ND	15	40	400	ND	ND	ND	1.0	ND
Males															
9–13 y	1,700	1,200	100	600	ND	ND	ND	20	60	600	ND	ND	ND	2.0	ND
14–18 y	2,800	1,800	100	800	ND	ND	ND	30	80	800	ND	ND	ND	3.0	ND
19–30 y	3,000	2,000	100	1,000	ND	ND	ND	35	100	1,000	ND	ND	ND	3.5	ND
31–50 y	3,000	2,000	100	1,000	ND	ND	ND	35	100	1,000	ND	ND	ND	3.5	ND
51–70 y	3,000	2,000	100	1,000	ND	ND	ND	35	100	1,000	ND	ND	ND	3.5	ND
> 70 y	3,000	2,000	100	1,000	ND	ND	ND	35	100	1,000	ND	ND	ND	3.5	ND
Females															
9–13 y	1,700	1,200	100	600	ND	ND	ND	20	60	600	ND	ND	ND	2.0	ND
14–18 y	2,800	1,800	100	800	ND	ND	ND	30	80	800	ND	ND	ND	3.0	ND
19–30 y	3,000	2,000	100	1,000	ND	ND	ND	35	100	1,000	ND	ND	ND	3.5	ND
31–50 y	3,000	2,000	100	1,000	ND	ND	ND	35	100	1,000	ND	ND	ND	3.5	ND
51–70 y	3,000	2,000	100	1,000	ND	ND	ND	35	100	1,000	ND	ND	ND	3.5	ND
> 70 y	3,000	2,000	100	1,000	ND	ND	ND	35	100	1,000	ND	ND	ND	3.5	ND
Pregnancy															
14–18 y	2,800	1,800	100	800	ND	ND	ND	30	80	800	ND	ND	ND	3.0	ND
19–30 y	3,000	2,000	100	1,000	ND	ND	ND	35	100	1,000	ND	ND	ND	3.5	ND
31–50 y	3,000	2,000	100	1,000	ND	ND	ND	35	100	1,000	ND	ND	ND	3.5	ND
Lactation															
14–18 y	2,800	1,800	100	800	ND	ND	ND	30	80	800	ND	ND	ND	3.0	ND
19–30 y	3,000	2,000	100	1,000	ND	ND	ND	35	100	1,000	ND	ND	ND	3.5	ND
31–50 y	3,000	2,000	100	1,000	ND	ND	ND	35	100	1,000	ND	ND	ND	3.5	ND

NOTE: A Tolerable Upper Intake Level (UL) is the highest level of daily nutrient intake that is likely to pose no risk of adverse health effects to almost all individuals in the general population. Unless otherwise specified, the UL represents total intake from food, water, and supplements. Due to a lack of suitable data, ULs could not be established for vitamin K, thiamin, riboflavin, vitamin B₁₂, pantothenic acid, biotin, and carotenoids. In the absence of a UL, extra caution may be warranted in consuming levels above recommended intakes. Members of the general population should be advised not to routinely exceed the UL. The UL is not meant to apply to individuals who are treated with the nutrient under medical supervision or to individuals with predisposing conditions that modify their sensitivity to the nutrient.

^aAs preformed vitamin A only.

^bAs α-tocopherol; applies to any form of supplemental α-tocopherol.

^cThe ULs for vitamin E, niacin, and folate apply to synthetic forms obtained from supplements, fortified foods, or a combination of the two.

^dβ-Carotene supplements are advised only to serve as a provitamin A source for individuals at risk of vitamin A deficiency.

^eND = Not determinable due to lack of data of adverse effects in this age group and concern with regard to lack of ability to handle excess amounts. Source of intake should be from food only to prevent high levels of intake.

SOURCES: *Dietary Reference Intakes for Calcium, Phosphorous, Magnesium, Vitamin D, and Fluoride* (1997); *Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B₆, Folate, Vitamin B₁₂, Pantothenic Acid, Biotin, and Choline* (1998); *Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium, and Carotenoids* (2000); *Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc* (2001); and *Dietary Reference Intakes for Calcium and Vitamin D* (2011). These reports may be accessed via www.nap.edu.

<http://www.nationalacademies.org/hmd/Activities/Nutrition/SummaryDRIs/DRI-Tables.aspx>

accessed 5/2/2017

Note page 2 for children 4-8 years - Tolerable Upper Intake Level for Sodium is 1.9 g/d = 1,900 mg/day

Dietary Reference Intakes (DRIs): Tolerable Upper Intake Levels, Elements
 Food and Nutrition Board, Institute of Medicine, National Academies

Life Stage Group	Arsenic ^a	Boron (mg/d)	Calcium (mg/d)	Chromium	Copper (μg/d)	Fluoride (mg/d)	Iodine (μg/d)	Iron (mg/d)	Magnesium (mg/d) ^b	Manganese (mg/d)	Molybdenum (μg/d)	Nickel (mg/d)	Phosphorus (g/d)	Selenium (μg/d)	Silicon ^c	Vanadium (mg/d) ^d	Zinc (mg/d)	Sodium (g/d)	Chloride (g/d)
Infants																			
0 to 6 mo	ND ^e	ND	1,000	ND	ND	0.7	ND	40	ND	ND	ND	ND	ND	45	ND	ND	4	ND	ND
6 to 12 mo	ND	ND	1,500	ND	ND	0.9	ND	40	ND	ND	ND	ND	ND	60	ND	ND	5	ND	ND
Children																			
1–3 y	ND	3	2,500	ND	1,000	1.3	200	40	65	2	300	0.2	3	90	ND	ND	7	1.5	2.3
4–8 y	ND	6	2,500	ND	3,000	2.2	300	40	110	3	600	0.3	3	150	ND	ND	12	1.9	2.9
Males																			
9–13 y	ND	11	3,000	ND	5,000	10	600	40	350	6	1,100	0.6	4	280	ND	ND	23	2.2	3.4
14–18 y	ND	17	3,000	ND	8,000	10	900	45	350	9	1,700	1.0	4	400	ND	ND	34	2.3	3.6
19–30 y	ND	20	2,500	ND	10,000	10	1,100	45	350	11	2,000	1.0	4	400	ND	1.8	40	2.3	3.6
31–50 y	ND	20	2,500	ND	10,000	10	1,100	45	350	11	2,000	1.0	4	400	ND	1.8	40	2.3	3.6
51–70 y	ND	20	2,000	ND	10,000	10	1,100	45	350	11	2,000	1.0	4	400	ND	1.8	40	2.3	3.6
> 70 y	ND	20	2,000	ND	10,000	10	1,100	45	350	11	2,000	1.0	3	400	ND	1.8	40	2.3	3.6
Females																			
9–13 y	ND	11	3,000	ND	5,000	10	600	40	350	6	1,100	0.6	4	280	ND	ND	23	2.2	3.4
14–18 y	ND	17	3,000	ND	8,000	10	900	45	350	9	1,700	1.0	4	400	ND	ND	34	2.3	3.6
19–30 y	ND	20	2,500	ND	10,000	10	1,100	45	350	11	2,000	1.0	4	400	ND	1.8	40	2.3	3.6
31–50 y	ND	20	2,500	ND	10,000	10	1,100	45	350	11	2,000	1.0	4	400	ND	1.8	40	2.3	3.6
51–70 y	ND	20	2,000	ND	10,000	10	1,100	45	350	11	2,000	1.0	4	400	ND	1.8	40	2.3	3.6
> 70 y	ND	20	2,000	ND	10,000	10	1,100	45	350	11	2,000	1.0	3	400	ND	1.8	40	2.3	3.6
Pregnancy																			
14–18 y	ND	17	3,000	ND	8,000	10	900	45	350	9	1,700	1.0	3.5	400	ND	ND	34	2.3	3.6
19–30 y	ND	20	2,500	ND	10,000	10	1,100	45	350	11	2,000	1.0	3.5	400	ND	ND	40	2.3	3.6
31–50 y	ND	20	2,500	ND	10,000	10	1,100	45	350	11	2,000	1.0	3.5	400	ND	ND	40	2.3	3.6
Lactation																			
14–18 y	ND	17	3,000	ND	8,000	10	900	45	350	9	1,700	1.0	4	400	ND	ND	34	2.3	3.6
19–30 y	ND	20	2,500	ND	10,000	10	1,100	45	350	11	2,000	1.0	4	400	ND	ND	40	2.3	3.6
31–50 y	ND	20	2,500	ND	10,000	10	1,100	45	350	11	2,000	1.0	4	400	ND	ND	40	2.3	3.6

NOTE: A Tolerable Upper Intake Level (UL) is the highest level of daily nutrient intake that is likely to pose no risk of adverse health effects to almost all individuals in the general population. Unless otherwise specified, the UL represents total intake from food, water, and supplements. Due to a lack of suitable data, ULs could not be established for vitamin K, thiamin, riboflavin, vitamin B₁₂, pantothenic acid, biotin, and carotenoids. In the absence of a UL, extra caution may be warranted in consuming levels above recommended intakes. Members of the general population should be advised not to routinely exceed the UL. The UL is not meant to apply to individuals who are treated with the nutrient under medical supervision or to individuals with predisposing conditions that modify their sensitivity to the nutrient.

^aAlthough the UL was not determined for arsenic, there is no justification for adding arsenic to food or supplements.
^bThe ULs for magnesium represent intake from a pharmacological agent only and do not include intake from food and water.
^cAlthough silicon has not been shown to cause adverse effects in humans, there is no justification for adding silicon to supplements.

^dAlthough vanadium in food has not been shown to cause adverse effects in humans, there is no justification for adding vanadium to food and vanadium supplements should be used with caution. The UL is based on adverse effects in laboratory animals and this data could be used to set a UL for adults but not children and adolescents.

^eND = Not determinable due to lack of data of adverse effects in this age group and concern with regard to lack of ability to handle excess amounts. Source of intake should be from food only to prevent high levels of intake.

SOURCES: *Dietary Reference Intakes for Calcium, Phosphorus, Magnesium, Vitamin D, and Fluoride* (1997); *Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B₆, Folate, Vitamin B₁₂, Pantothenic Acid, Biotin, and Choline* (1998); *Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium, and Carotenoids* (2000); *Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc* (2001); *Dietary Reference Intakes for Water, Potassium, Sodium, Chloride, and Sulfate* (2005); and *Dietary Reference Intakes for Calcium and Vitamin D* (2011). These reports may be accessed via www.nap.edu.

<http://www.nationalacademies.org/hmd/Activities/Nutrition/SummaryDRIs/DRI-Tables.aspx>

accessed 5/2/2017